

Affine parts of abelian surfaces as complete intersection of three quartics

A. Lesfari

Department of Mathematics

Faculty of Sciences

University of Chouaib Doukkali

B.P. 20, El-Jadida, Morocco.

E. mail : Lesfariahmed@yahoo.fr, Lesfari@ucd.ac.ma

Abstract

We consider an integrable system in five unknowns having three quartics invariants. We show that the complex affine variety defined by putting these invariants equal to generic constants, completes into an abelian surface; the jacobian of a genus two hyperelliptic curve. This system is algebraic completely integrable and it can be integrated in genus two hyperelliptic functions.

Keywords. Integrable systems, curves, Kummer surfaces, Abelian surfaces.

Mathematics Subject Classification (2000). 70H06, 37J35, 14H70, 14K20, 14H40.

1 Introduction

The problem of finding and integrating hamiltonian systems, has attracted a considerable amount of attention in recent years. Beside the fact that many integrable hamiltonian systems have been on the subject of powerful and beautiful theories of mathematics, another motivation for its study is: the concepts of integrability have been applied to an increasing number of physical systems, biological phenomena, population dynamics, chemical rate equations, to mention only a few. However, it seems still hopeless to describe or even to recognize with any facility, those hamiltonian systems which are integrable, though they are quite exceptional. In this paper, we shall be concerned with finite dimensional algebraic completely integrable systems. A dynamical system is algebraic completely integrable if it can be linearized on a complex algebraic torus $\mathbb{C}^n/lattice$ (=abelian variety). The invariants (often called first integrals or constants) of the motion are polynomials and the phase space coordinates (or some algebraic functions of these) restricted

to a complex invariant variety defined by putting these invariants equals to generic constants, are meromorphic functions on an abelian variety. Moreover, in the coordinates of this abelian variety, the flows (run with complex time) generated by the constants of the motion are straight lines. However, besides the fact that many hamiltonian completely integrable systems possess this structure, another motivation for its study which sounds more modern is: algebraic completely integrable systems come up systematically whenever you study the isospectral deformation of some linear operator containing a rational indeterminate. Therefore there are hidden symmetries which have a group theoretical foundation. The concept of algebraic complete integrability is quite effective in small dimensions and has the advantage to lead to global results, unlike the existing criteria for real analytic integrability, which, at this stage are perturbation results. In fact, the overwhelming majority of dynamical systems, hamiltonian or not, are non-integrable and possess regimes of chaotic behavior in phase space. In the present paper, we discuss an interesting interaction between complex algebraic geometry and dynamical systems. we present an integrable system in five unknowns having three quartics invariants. This system is algebraic completely integrable in \mathbb{C}^5 , it can be integrated in genus 2 hyperelliptic functions. We show that the complex affine variety $B(3)$ defined by putting these invariants equal to generic constants, is a double cover of a Kummer surface and the system (1) can be integrated in genus 2 hyperelliptic functions. We make a careful study of the algebraic geometric aspect of the affine variety $B(3)$ of the system (1). We find via the Painlevé analysis the principal balances of the hamiltonian field defined by the hamiltonian. To be more precise, we show that the system (1) possesses Laurent series solutions in t , which depend on 4 free parameters : α, β, γ and θ . These meromorphic solutions restricted to the surface $B(3)$ are parameterized by two isomorphic smooth hyperelliptic curves $\mathcal{H}_{\varepsilon=\pm i}(8)$ of genus 2 that intersect in only one point at which they are tangent to each other. The affine variety $B(3)$ is embedded into \mathbb{P}^{15} and completes into an abelian variety \tilde{B} (the jacobian of a genus 2 curve) by adjoining a divisor $\mathcal{D} = \mathcal{H}_i + \mathcal{H}_{-i}$. The latter has geometric genus 5 and $\mathcal{S} = 2\mathcal{D}$ (very ample) has genus 17. The flow (1) evolves on \tilde{B} and is tangent to each hyperelliptic curve \mathcal{H}_ε at the point of tangency between them. Consequently, the system (1) is algebraic integrable.

Abelian varieties, very heavily studied by algebraic geometers, enjoy certain algebraic properties which can then be translated into differential equations and their Laurent solutions. Among the results presented in this paper, there is an explicit calculation of invariants for a hamiltonian system which cut out an open set in an abelian variety and various curves related to this system are given explicitly. The integrable dynamical system presented here is interesting, particular to experts of abelian varieties who may want to see explicit examples of a correspondence for varieties defined by different curves.

2 A five-dimensional integrable system

Let us consider the following system of five differential equations in the unknowns z_1, \dots, z_5 :

$$\begin{aligned}
 \dot{z}_1 &= 2z_4, \\
 \dot{z}_2 &= z_3, \\
 \dot{z}_3 &= -4az_2 - 6z_1z_2 - 16z_2^3, \\
 \dot{z}_4 &= -az_1 - z_1^2 - 8z_1z_2^2 + z_5, \\
 \dot{z}_5 &= -8z_2^2z_4 - 2az_4 - 2z_1z_4 + 4z_1z_2z_3,
 \end{aligned} \tag{1}$$

where the dot denotes differentiation with respect to the time t .

Proposition 2.1 *The system (1) possesses three quartic invariants and is completely integrable in the sense of Liouville. The complex affine variety $B(3)$ defined by putting these invariants equal to generic constants, is a double cover of a Kummer surface(4) and the system (1) can be integrated in genus 2 hyperelliptic functions.*

Proof. The following three quartics are constants of motion for this system

$$\begin{aligned}
 F_1 &= \frac{1}{2}z_5 + 2z_1z_2^2 + \frac{1}{2}z_3^2 + \frac{1}{2}az_1 + 2az_2^2 + \frac{1}{4}z_1^2 + 4z_2^4, \\
 F_2 &= az_1z_2 + z_1^2z_2 + 4z_1z_2^3 - z_2z_5 + z_3z_4, \\
 F_3 &= z_1z_5 - 2z_1^2z_2^2 - z_4^2.
 \end{aligned} \tag{2}$$

The system (1) can be written as a hamiltonian vector field

$$\dot{z} = J \frac{\partial H}{\partial z}, \quad z = (z_1, z_2, z_3, z_4, z_5)^\top,$$

where $H = F_1$. The hamiltonian structure is defined by the Poisson bracket

$$\{F, H\} = \left\langle \frac{\partial F}{\partial z}, J \frac{\partial H}{\partial z} \right\rangle = \sum_{k,l=1}^5 J_{kl} \frac{\partial F}{\partial z_k} \frac{\partial H}{\partial z_l},$$

where

$$\frac{\partial H}{\partial z} = \left(\frac{\partial H}{\partial z_1}, \frac{\partial H}{\partial z_2}, \frac{\partial H}{\partial z_3}, \frac{\partial H}{\partial z_4}, \frac{\partial H}{\partial z_5} \right)^\top,$$

and

$$J = \begin{bmatrix} 0 & 0 & 0 & 2z_1 & 4z_4 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & -1 & 0 & 0 & -4z_1z_2 \\ -2z_1 & 0 & 0 & 0 & 2z_5 - 8z_1z_2^2 \\ -4z_4 & 0 & 4z_1z_2 & -2z_5 + 8z_1z_2^2 & 0 \end{bmatrix},$$

is a skew-symmetric matrix for which the corresponding Poisson bracket satisfies the Jacobi identities. The second flow commuting with the first is regulated by the equations

$$\dot{z} = J \frac{\partial F_2}{\partial z}, \quad z = (z_1, z_2, z_3, z_4, z_5)^\top,$$

and is written explicitly as

$$\begin{aligned} \dot{z}_1 &= 2z_1z_3 - 4z_2z_4, \\ \dot{z}_2 &= z_4, \\ \dot{z}_3 &= z_5 - 8z_1z_2^2 - az_1 - z_1^2, \\ \dot{z}_4 &= -2az_1z_2 - 4z_1^2z_2 - 2z_2z_5, \\ \dot{z}_5 &= -4az_2z_4 - 4z_1z_2z_4 - 16z_2^3z_4 - 2z_3z_5 + 8z_1z_2^2z_3. \end{aligned}$$

These vector fields are in involution, i.e.,

$$\{F_1, F_2\} = \left\langle \frac{\partial F_1}{\partial z}, J \frac{\partial F_2}{\partial z} \right\rangle = 0,$$

and the remaining one is casimir, i.e.,

$$J \frac{\partial F_3}{\partial z} = 0.$$

Let B be the complex affine variety defined by

$$B = \bigcap_{k=1}^2 \{z : F_k(z) = c_k\} \subset \mathbb{C}^5. \quad (3)$$

Since B is the fibre of a morphism from \mathbb{C}^5 to \mathbb{C}^3 over $(c_1, c_2, c_3) \in \mathbb{C}^3$, for almost all c_1, c_2, c_3 , therefore B is a smooth affine surface. Note that

$$\sigma : (z_1, z_2, z_3, z_4, z_5) \mapsto (z_1, z_2, -z_3, -z_4, z_5),$$

is an involution on B . The quotient B/σ is a Kummer surface defined by

$$p(z_1, z_2)z_5^2 + q(z_1, z_2)z_5 + r(z_1, z_2) = 0, \quad (4)$$

where

$$\begin{aligned} p(z_1, z_2) &= z_2^2 + z_1, \\ q(z_1, z_2) &= \frac{1}{2}z_1^3 + 2az_1z_2^2 + az_1^2 - 2c_1z_1 + 2c_2z_2 - c_3, \\ r(z_1, z_2) &= -8c_3z_2^4 + (a^2 + 4c_1)z_1^2z_2^2 - 8c_2z_1z_2^3 - 2c_2z_1^2z_2 - 4c_3z_1z_2^2 \\ &\quad - \frac{1}{2}c_3z_1^2 - 4ac_3z_2^2 - 2ac_2z_1z_2 - ac_3z_1 + c_2^2 + 2c_1c_3. \end{aligned}$$

Using $F_1 = c_1$ (2), we have

$$z_5 = 2c_1 - 4z_1z_2^2 - z_3^2 - az_1 - 4az_2^2 - \frac{1}{2}z_1^2 - 8z_2^4,$$

and substituting this into $F_2 = c_2$, $F_3 = c_3$, (2) yields

$$\begin{aligned} 2az_1z_2 + \frac{3}{2}z_1^2z_2 + 8z_1z_2^3 - 2c_1z_2 + z_2z_3^2 + 4az_2^3 + 8z_2^5 + z_3z_4 &= c_2, \\ 2c_1z_1 - 6z_1^2z_2^2 - z_1z_3^2 - az_1^2 - 4az_1z_2^2 - \frac{1}{2}z_1^3 - 8z_1z_2^4 - z_4^2 &= c_3. \end{aligned} \quad (5)$$

We introduce two coordinates s_1, s_2 as follows

$$\begin{aligned} z_1 &= -4s_1s_2, \\ z_2 &= s_1 + s_2, \\ z_3 &= \dot{s}_1 + \dot{s}_2, \\ z_4 &= -2(\dot{s}_1s_2 + s_1\dot{s}_2). \end{aligned}$$

Upon substituting this parametrization, (5) turns into

$$\begin{aligned} (s_1 - s_2) ((\dot{s}_1)^2 - (\dot{s}_2)^2) + 8(s_1 + s_2)(s_1^4 + s_2^4 + s_1^2s_2^2) \\ + 4a(s_1 + s_2)(s_1^2 + s_2^2) - 2c_1(s_1 + s_2) - c_2 = 0, \\ (s_1 - s_2)(s_2(\dot{s}_1)^2 - s_1(\dot{s}_2)^2) + 32s_1s_2(s_1^4 + s_2^4 + s_1^2s_2^2) \\ + 32s_1^2s_2^2(s_1^2 + s_2^2) + 16as_1s_2(s_1^2 + s_2^2) + 16as_1^2s_2^2 - 8c_1s_1s_2 - c_3 = 0. \end{aligned}$$

These equations are solved linearly for \dot{s}_1^2 and \dot{s}_2^2 as

$$\begin{aligned} (\dot{s}_1)^2 &= \frac{-32s_1^6 - 16as_1^4 + 8c_1s_1^2 + 4c_2s_1 - c_3}{4(s_2 - s_1)^2}, \\ (\dot{s}_2)^2 &= \frac{-32s_2^6 - 16as_2^4 + 8c_1s_2^2 + 4c_2s_2 - c_3}{4(s_2 - s_1)^2}, \end{aligned} \quad (6)$$

and can be integrated by means of the Abel transformation $\mathcal{H} \rightarrow \text{Jac}(\mathcal{H})$, where the hyperelliptic curve \mathcal{H} of genus 2 is given by an equation

$$w^2 = -32s^6 - 16as^4 + 8c_1s^2 + 4c_2s - c_3.$$

Consequently, the equations (1) are integrated in terms of genus 2 hyperelliptic functions. This establishes the proposition.

3 Laurent series solutions and algebraic curves

The invariant variety $B(3)$ is a smooth affine surface for generic values of c_1, c_2 and c_3 . So, the question I address is how does one find the compactification of B into an abelian surface? Following the methods in Adler-van

Moerbeke [1], the idea of the direct proof is closely related to the geometric spirit of the (real) Arnold-Liouville theorem [8]. Namely, a compact complex n -dimensional variety on which there exist n holomorphic commuting vector fields which are independent at every point is analytically isomorphic to a n -dimensional complex torus $\mathbb{C}^n/Lattice$ and the complex flows generated by the vector fields are straight lines on this complex torus. Now, the main problem will be to complete $B(3)$ into a non singular compact complex algebraic variety $\tilde{B} = B \cup \mathcal{D}$ in such a way that the vector fields X_{F_1} and X_{F_2} generated respectively by F_1 and F_2 , extend holomorphically along a divisor \mathcal{D} and remain independent there. If this is possible, \tilde{B} is an algebraic complex torus (an abelian variety) and the coordinates z_1, \dots, z_5 restricted to B are abelian functions. A naive guess would be to take the natural compactification \overline{B} of B by projectivizing the equations:

$$\overline{B} = \bigcap_{k=1}^3 \{F_k(Z) = c_k Z_0^4\} \subset \mathbb{P}^5.$$

Indeed, this can never work for a general reason : an abelian variety \tilde{B} of dimension bigger or equal than two is never a complete smooth intersection, that is it can never be described in some projective space \mathbb{P}^n by $n - \dim \tilde{B}$ global polynomial homogeneous equations. In other words, if B is to be the affine part of an abelian surface, \overline{B} must have a singularity somewhere along the locus at infinity $\overline{B} \cap \{Z_0 = 0\}$. In fact, we shall show that the existence of meromorphic solutions to the differential equations (1) depending on 4 free parameters can be used to manufacture the tori, without ever going through the delicate procedure of blowing up and down. Information about the tori can then be gathered from the divisor.

Proposition 3.1 *The system (1) possesses Laurent series solutions which depend on 4 free parameters : α, β, γ and θ . These meromorphic solutions restricted to the surface $B(3)$ are parameterized by two isomorphic smooth hyperelliptic curves $\mathcal{H}_{\varepsilon=\pm i}(8)$ of genus 2.*

Proof. The first fact to observe is that if the system is to have Laurent solutions depending on 4 free parameters, the Laurent decomposition of such asymptotic solutions must have the following form

$$\begin{aligned} z_1 &= \frac{1}{t}(z_1^{(0)} + z_1^{(1)}t + z_1^{(2)}t^2 + z_1^{(3)}t^3 + z_1^{(4)}t^4 + \dots), \\ z_2 &= \frac{1}{t}(z_2^{(0)} + z_2^{(1)}t + z_2^{(2)}t^2 + z_2^{(3)}t^3 + z_2^{(4)}t^4 + \dots), \\ z_3 &= \frac{1}{t^2}(-z_2^{(0)} + z_2^{(2)}t^2 + 2z_2^{(3)}t^3 + 3z_2^{(4)}t^4 + \dots), \\ z_4 &= \frac{1}{2t^2}(-z_1^{(0)} + z_1^{(2)}t^2 + 2z_1^{(3t)}t^3 + 3z_1^{(4)}t^4 + \dots), \\ z_5 &= \frac{1}{t^3}(z_5^{(0)} + z_5^{(1)}t + z_5^{(2)}t^2 + z_5^{(3)}t^3 + z_5^{(4)}t^4 + \dots). \end{aligned}$$

Putting these expansions into

$$\begin{aligned}\ddot{z}_1 &= -2az_1 - 2z_1^2 - 16z_1z_2^2 + 2z_5, \\ \ddot{z}_2 &= -4az_2 - 6z_1z_2 - 16z_2^3, \\ \ddot{z}_5 &= -8z_2^2z_4 - 2az_4 - 2z_1z_4 + 4z_1z_2z_3,\end{aligned}$$

deduced from (1), solving inductively for the $z_k^{(j)}$ ($k = 1, 2, 5$), one finds at the 0^{th} step (resp. 2^{th} step) a free parameter α (resp. β) and the two remaining ones γ, θ at the 4^{th} step. More precisely, we have

$$\begin{aligned}z_1 &= \frac{1}{t}(\alpha - \alpha^2t + \beta t^2 + \frac{1}{6}\alpha(3\beta - 9\alpha^3 + 4a\alpha)t^3 + \gamma t^4 + \dots), \\ z_2 &= \frac{\varepsilon\sqrt{2}}{4t}(1 + \alpha t + \frac{1}{3}(-3\alpha^2 + 2a)t^2 + \frac{1}{2}(3\beta - \alpha^3)t^3 - 2\varepsilon\sqrt{2}\theta t^4 + \dots), \\ z_3 &= \frac{\varepsilon\sqrt{2}}{4t^2}(-1 + \frac{1}{3}(-3\alpha^2 + 2a)t^2 + (3\beta - \alpha^3)t^3 - 6\varepsilon\sqrt{2}\theta t^4 + \dots), \\ z_4 &= \frac{1}{2t^2}(-\alpha + \beta t^2 + \frac{1}{3}\alpha(3\beta - 9\alpha^3 + 4a\alpha)t^3 + 3\gamma t^4 + \dots), \\ z_5 &= \frac{1}{t}(-\frac{1}{3}a\alpha + \alpha^3 - \beta + (3\alpha^4 - a\alpha^2 - 3\alpha\beta)t \\ &\quad + (4\varepsilon\sqrt{2}\alpha\theta + 2\gamma + \frac{8}{3}a\alpha^3 - \frac{1}{3}a\beta - \alpha^2\beta - 3\alpha^5 - \frac{4}{9}a^2\alpha)t^2 + \dots),\end{aligned}\tag{7}$$

with $\varepsilon = \pm i$. Using the majorant method, we can show that the formal Laurent series solutions are convergent. Substituting the solutions (7) into $F_1 = c_1$, $F_2 = c_2$ and $F_3 = c_3$, and equating the t^0 -terms yields

$$\begin{aligned}F_1 &= \frac{15}{8}\alpha^4 - \frac{5}{6}a\alpha^2 - \frac{5}{4}\alpha\beta - \frac{7}{36}a^2 - \frac{5}{4}\varepsilon\sqrt{2}\theta = c_1, \\ F_2 &= \varepsilon\sqrt{2}(\frac{1}{4}\alpha^5 - \gamma + \frac{\varepsilon\sqrt{2}}{2}\alpha\theta - \frac{2}{3}a\alpha^3 + \frac{1}{3}a\beta + \frac{1}{6}a^2 + \frac{1}{2}\alpha^2\beta) = c_2, \\ F_3 &= -\frac{11}{2}\alpha^6 - \beta^2 + 4a\gamma + 3\alpha^2\varepsilon\sqrt{2}\theta + \alpha^3\beta - \frac{1}{3}a^2\alpha^2 + \frac{10}{3}a\alpha^4 = c_3.\end{aligned}$$

Eliminating γ and θ from these equations, leads to an equation connecting the two remaining parameters α and β :

$$\beta^2 + \frac{2}{3}(3\alpha^2 - 2a)\alpha\beta - 3\alpha^6 + \frac{8}{3}a\alpha^4 + \frac{4}{9}(a^2 + 9c_1)\alpha^2 - 2\varepsilon\sqrt{2}c_2\alpha + c_3 = 0, \tag{8}$$

According to Hurwitz' formula, this defines two isomorphic smooth hyperelliptic curves \mathcal{H}_ε ($\varepsilon = \pm i$) of genus 2, which finishes the proof of the proposition.

4 Affine part of an abelian surface as the jacobian of a genus two hyperelliptic curve

In order to embed \mathcal{H}_ε into some projective space, one of the key underlying principles used is the Kodaira embedding theorem, which states that a

smooth complex manifold can be smoothly embedded into projective space \mathbb{P}^N with the set of functions having a pole of order k along positive divisor on the manifold, provided k is large enough; fortunately, for abelian varieties, k need not be larger than three according to Lefschetz. These functions are easily constructed from the Laurent solutions (7) by looking for polynomials in the phase variables which in the expansions have at most a k -fold pole. The nature of the expansions and some algebraic proprieties of abelian varieties provide a recipe for when to terminate our search for such functions, thus making the procedure implementable. Precisely, we wish to find a set of polynomial functions $\{f_0, \dots, f_N\}$, of increasing degree in the original variables z_1, \dots, z_5 , having the property that the embedding \mathcal{D} of $\mathcal{H}_i + \mathcal{H}_{-i}$ into \mathbb{P}^N via those functions satisfies the relation :

$$\text{geometric genus}(2\mathcal{D}) \equiv g(2\mathcal{D}) = N + 2.$$

A this point, it may be not so clear why the curve \mathcal{D} must really live on an abelian surface. Let us say, for the moment, that the equations of the divisor \mathcal{D} (i.e., the place where the solutions blow up), as a curve traced on the abelian surface \tilde{B} (to be constructed in proposition 4.2), must be understood as relations connecting the free parameters as they appear firstly in the expansions (7). In the present situation, this means that (8) must be understood as relations connecting α and β . Let

$$L^{(r)} = \left\{ \begin{array}{l} \text{polynomials } f = f(z, \dots, z_5) \\ \text{of degree } \leq r, \text{ with at worst a} \\ \text{double pole along } \mathcal{H}_i + \mathcal{H}_{-i} \\ \text{and with } z_1, \dots, z_5 \text{ as in(7)} \end{array} \right\} / [F_k = c_k, k = 1, 2, 3],$$

and let $(f_0, f_1, \dots, f_{N_r})$ be a basis of $L^{(r)}$. We look for r such that :

$$g(2\mathcal{D}^{(r)}) = N_r + 2, \quad 2\mathcal{D}^{(r)} \subset \mathbb{P}^{N_r}.$$

We shall show (proposition 4.1) that it is unnecessary to go beyond $r=4$.

Lemma 4.1 *The spaces $L^{(r)}$, nested according to weighted degree, are generated as follows*

$$\begin{aligned} L^{(1)} &= \{f_0, f_1, f_2, f_3, f_4, f_5\}, \\ L^{(2)} &= L^{(1)} \oplus \{f_6, f_8, f_9, f_{10}, f_{11}, f_{12}\}, \\ L^{(3)} &= L^{(2)}, \\ L^{(4)} &= L^{(3)} \oplus \{f_{13}, f_{14}, f_{15}\}, \end{aligned} \tag{9}$$

where

$$f_0 = 1, \quad f_1 = z_1 = \frac{\alpha}{t} + \dots,$$

$$\begin{aligned}
f_2 = z_2 &= \frac{\varepsilon\sqrt{2}}{t} + \dots, & f_3 = z_3 &= -\frac{\varepsilon\sqrt{2}}{4t^2} + \dots, \\
f_4 = z_4 &= -\frac{\alpha}{2t^2} + \dots, & f_5 = z_5 &= -\frac{\eta}{3t} + \dots, \\
f_6 = z_1^2 &= \frac{\alpha^2}{t^2} + \dots, & f_7 = z_2^2 &= -\frac{1}{8t^2} + \dots, \\
f_8 = z_5^2 &= \frac{\eta^2}{9t^2} + \dots, & f_9 = z_1 z_2 &= \frac{\varepsilon\sqrt{2}\alpha}{4t^2} + \dots, \\
f_{10} = z_1 z_5 &= -\frac{\alpha\eta}{3t^2} + \dots, & f_{11} = z_2 z_5 &= -\frac{\varepsilon\sqrt{2}\eta}{12t^2} + \dots, \\
f_{12} = [z_1, z_2] &= -\frac{\varepsilon\sqrt{2}\alpha^2}{2t^2} + \dots, & f_{13} = [z_1, z_5] &= \frac{4\alpha^2\eta}{3t^2} + \dots, \\
f_{14} = [z_2, z_5] &= \frac{\varepsilon\sqrt{2}\alpha\eta}{6t^2} + \dots, & f_{15} = (z_3 - 2\varepsilon\sqrt{2}z_2^2)^2 &= -\frac{\alpha^2}{2t^2} + \dots,
\end{aligned}$$

with

$$[z_j, z_k] = \dot{z}_j z_k - z_j \dot{z}_k,$$

the wronskien of z_k and z_j , and

$$\eta \equiv 3\beta - 3\alpha^3 + \alpha\alpha.$$

Proof. The proof of this lemma is straightforward and can be done by inspection of the expansions (7). Note also that the functions z_1, z_2, z_5 behave as $1/t$ and if we consider the derivatives of the ratios $z_1/z_2, z_1/z_5, z_2/z_5$, the wronskiens $[z_1, z_2], [z_1, z_5], [z_2, z_5]$, must behave as $1/t^2$ since z_2^2, z_5^2 behave as $1/t^2$. This finishes the proof of the lemma.

Note that

$$\dim L^{(1)} = 6, \quad \dim L^{(2)} = \dim L^{(3)} = 13, \quad \dim L^{(4)} = 16.$$

Proposition 4.1 $L^{(4)}$ provides an embedding of $\mathcal{D}^{(4)}$ into projective space \mathbb{P}^{15} and $\mathcal{D}^{(4)}$ (resp. $2\mathcal{D}^{(4)}$) has genus 5 (resp. 17).

Proof. It turns out that neither $L^{(1)}$, nor $L^{(2)}$, nor $L^{(3)}$, yields a curve of the right genus; in fact

$$g(2\mathcal{D}^{(r)}) \neq \dim L^{(r)} + 1, \quad r = 1, 2, 3.$$

For instance, the embedding into \mathbb{P}^5 via $L^{(1)}$ does not separate the sheets, so we proceed to $L^{(2)}$ and we consider the corresponding embedding into \mathbb{P}^{12} . For finite values of α and β , the curves \mathcal{H}_i and \mathcal{H}_{-i} are disjoint; dividing the vector (f_0, \dots, f_{12}) by f_7 and taking the limit $t \rightarrow 0$, to yield

$$[0 : 0 : 0 : 2\varepsilon\sqrt{2} : 4\alpha : 0 : -8\alpha^2 : 1 : -\frac{8}{9}\eta^2 : -2\varepsilon\sqrt{2}\alpha : \frac{8}{3}\alpha\eta : \frac{2\varepsilon\sqrt{2}}{3}\eta : 4\varepsilon\sqrt{2}\alpha^2].$$

The curve (8) has two points covering $\alpha = \infty$, at which $\eta \equiv 3\beta - 3\alpha^3 + a\alpha$ behaves as follows :

$$\begin{aligned}\eta &= -6\alpha^3 + 3a\alpha \pm 3\sqrt{4\alpha^6 - 4a\alpha^4 - 4c_1\alpha^2 + 2\varepsilon\sqrt{2}c_2\alpha - c_3}, \\ &= \begin{cases} -\frac{3(a^2+4c_1)}{4\alpha} + & \text{lower order terms, picking the + sign,} \\ -12\alpha^3 + & O(\alpha), \text{ picking the - sign.} \end{cases}\end{aligned}$$

Then by picking the - sign and by dividing the vector (f_0, \dots, f_{12}) by f_8 , the corresponding point is mapped into the point

$$[0 : 0 : 0 : 0 : 0 : 0 : 0 : 0 : 1 : 0 : 0 : 0 : 0],$$

in \mathbb{P}^{12} which is independent of ε , whereas picking the + sign leads to two different points, according to the sign of ε . Thus, adding at least 2 to the genus of each curve, so that

$$g(2\mathcal{D}^{(2)}) - 2 > 12, \quad 2\mathcal{D}^{(2)} \subset \mathbb{P}^{12} \neq \mathbb{P}^{g-2},$$

which contradicts the fact that

$$N_r = g(2\mathcal{D}^{(2)}) - 2.$$

The embedding via $L^{(2)}$ (or $L^{(3)}$) is unacceptable as well. Consider now the embedding $2\mathcal{D}^{(4)}$ into \mathbb{P}^{15} using the 16 functions f_0, \dots, f_{15} of $L^{(4)}(9)$. It is easily seen that these functions separate all points of the curve (except perhaps for the points at ∞) : The curves \mathcal{H}_i and \mathcal{H}_{-i} are disjoint for finite values of α and β ; dividing the vector (f_0, \dots, f_{15}) by f_7 and taking the limit $t \rightarrow 0$, to yield

$$\begin{aligned}[0 : 0 : 0 : 2\varepsilon\sqrt{2} : 4\alpha : 0 : -8\alpha^2 : 1 : -\frac{8}{9}\eta^2 : -2\varepsilon\sqrt{2}\alpha : \frac{8}{3}\alpha\eta : \frac{2\varepsilon\sqrt{2}}{3}\eta : \\ 4\varepsilon\sqrt{2}\alpha^2 : -\frac{32}{3}\alpha^2\eta : -\frac{4\varepsilon\sqrt{2}}{3}\alpha\eta : 4\alpha^2].\end{aligned}$$

About the point $\alpha = \infty$, it is appropriate to divide by g_8 ; then by picking the sign - in η above, the corresponding point is mapped into the point

$$[0 : 0 : 0 : 0 : 0 : 0 : 0 : 0 : 1 : 0 : 0 : 0 : 0 : 0 : 0],$$

in \mathbb{P}^{15} which is independent of ε , whereas picking the + sign leads to two different points, according to the sign of ε . Hence, the divisor $\mathcal{D}^{(4)}$ obtained in this way has genus 5 and thus $g(2\mathcal{D}^{(4)})$ has genus 17 and

$$2\mathcal{D}^{(4)} \subset \mathbb{P}^{15} = \mathbb{P}^{g-2},$$

as desired. This ends the proof of the proposition.

Let

$$L = L^{(4)}, \quad \mathcal{D} = \mathcal{D}^{(4)},$$

and

$$\mathcal{S} = 2\mathcal{D}^{(4)} \subset \mathbb{P}^{15}.$$

Next we wish to construct a surface strip around \mathcal{S} which will support the commuting vector fields. In fact, \mathcal{S} has a good chance to be very ample divisor on an abelian surface, still to be constructed.

Proposition 4.2 *The variety B (3) generically is the affine part of an abelian surface \tilde{B} , more precisely the jacobian of a genus 2 curve. The reduced divisor at infinity*

$$\tilde{B} \setminus B = \mathcal{H}_i + \mathcal{H}_{-i},$$

consists of two smooth isomorphic genus 2 curves $\mathcal{H}_\varepsilon(8)$. The system of differential equations (1) is algebraic complete integrable and the corresponding flows evolve on \tilde{B} .

Proof. We need to attach the affine part of the intersection of the three invariants (2) so as to obtain a smooth compact connected surface in \mathbb{P}^{15} . To be precise, the orbits of the vector field (1) running through \mathcal{S} form a smooth surface Σ near \mathcal{S} such that

$$\Sigma \setminus B \subseteq \tilde{B},$$

and the variety

$$\tilde{B} = B \cup \Sigma,$$

is smooth, compact and connected. Indeed, let

$$\psi(t, p) = \{z(t) = (z_1(t), \dots, z_5(t)) : t \in \mathbb{C}, 0 < |t| < \varepsilon\},$$

be the orbit of the vector field (1) going through the point $p \in \mathcal{S}$. Let $\Sigma_p \subset \mathbb{P}^{15}$ be the surface element formed by the divisor \mathcal{S} and the orbits going through p . Consider the curve

$$\mathcal{S}' = \mathcal{H} \cap \Sigma,$$

where $\mathcal{H} \subset \mathbb{P}^{15}$ is a hyperplane transversal to the direction of the flow and $\Sigma \equiv \cup_{p \in \mathcal{S}} \Sigma_p$. If \mathcal{S}' is smooth, then using the implicit function theorem the surface Σ is smooth. But if \mathcal{S}' is singular at 0, then Σ would be singular along the trajectory (t -axis) which go immediately into the affine part B . Hence, B would be singular which is a contradiction because B is the fibre of a morphism from \mathbb{C}^5 to \mathbb{C}^2 and so smooth for almost all the three constants of the motion c_k . Next, let \overline{B} be the projective closure of B into \mathbb{P}^5 , let

$Z = [Z_0 : Z_1 : \dots : Z_5] \in \mathbb{P}^5$ and let $I = \overline{B} \cap \{Z_0 = 0\}$ be the locus at infinity. Consider the map

$$\overline{B} \subseteq \mathbb{P}^5 \longrightarrow \mathbb{P}^{15}, \quad Z \longmapsto f(Z),$$

where $f = (f_0, f_1, \dots, f_{15}) \in L(\mathcal{S})(9)$ and let $\tilde{B} = f(\overline{B})$. In a neighbourhood $V(p) \subseteq \mathbb{P}^{15}$ of p , we have $\Sigma_p = \tilde{B}$ and $\Sigma_p \setminus \mathcal{S} \subseteq \tilde{B}$. Otherwise there would exist an element of surface $\Sigma'_p \subseteq \tilde{B}$ such that

$$\Sigma_p \cap \Sigma'_p = (t - axis), \text{ orbit } \psi(t, p) = (t - axis) \setminus p \subseteq \tilde{B},$$

and hence \tilde{B} would be singular along the t -axis which is impossible. Since the variety $\overline{B} \cap \{Z_0 \neq 0\}$ is irreducible and since the generic hyperplane section $\mathcal{H}_{gen.}$ of \overline{B} is also irreducible, all hyperplane sections are connected and hence I is also connected. Now, consider the graph $\Gamma_f \subseteq \mathbb{P}^5 \times \mathbb{P}^{15}$ of the map f , which is irreducible together with \overline{B} . It follows from the irreducibility of I that a generic hyperplane section $\Gamma_f \cap \{\mathcal{H}_{gen.} \times \mathbb{P}^{15}\}$ is irreducible, hence the special hyperplane section $\Gamma_f \cap \{\{Z_0 = 0\} \times \mathbb{P}^{15}\}$ is connected and therefore the projection map

$$proj_{\mathbb{P}^{15}} \{\Gamma_f \cap \{\{Z_0 = 0\} \times \mathbb{P}^{15}\}\} = f(I) \equiv \mathcal{S},$$

is connected. Hence, the variety $B \cup \Sigma = \tilde{B}$ is compact, connected and embeds smoothly into \mathbb{P}^{15} via f . We wish to show that \tilde{B} is an abelian surface equipped with two everywhere independent commuting vector fields. For doing that, let ϕ^{τ_1} and ϕ^{τ_2} be the flows corresponding to vector fields X_{F_1} and X_{F_2} . The latter are generated respectively by F_1 and F_2 . For $p \in \mathcal{S}$ and for small $\varepsilon > 0$,

$$\phi^{\tau_1}(p), \quad \forall \tau_1, 0 < |\tau_1| < \varepsilon,$$

is well defined and $\phi^{\tau_1}(p) \in B$. Then we may define ϕ^{τ_2} on B by

$$\phi^{\tau_2}(q) = \phi^{-\tau_1} \phi^{\tau_2} \phi^{\tau_1}(q), \quad q \in U(p) = \phi^{-\tau_1}(U(\phi^{\tau_1}(p))),$$

where $U(p)$ is a neighbourhood of p . By commutativity one can see that ϕ^{τ_2} is independent of τ_1 ;

$$\begin{aligned} \phi^{-\tau_1 - \varepsilon_1} \phi^{\tau_2} \phi^{\tau_1 + \varepsilon_1}(q) &= \phi^{-\tau_1} \phi^{-\varepsilon_1} \phi^{\tau_2} \phi^{\tau_1} \phi^{\varepsilon_1}, \\ &= \phi^{-\tau_1} \phi^{\tau_2} \phi^{\tau_1}(q). \end{aligned}$$

We affirm that $\phi^{\tau_2}(q)$ is holomorphic away from \mathcal{S} . This because $\phi^{\tau_2} \phi^{\tau_1}(q)$ is holomorphic away from \mathcal{S} and that ϕ^{τ_1} is holomorphic in $U(p)$ and maps bi-holomorphically $U(p)$ onto $U(\phi^{\tau_1}(p))$. Now, since the flows ϕ^{τ_1} and ϕ^{τ_2} are holomorphic and independent on \mathcal{S} , we can show along the same lines as in the Arnold-Liouville theorem [1,6] that \tilde{B} is a complex torus $\mathbb{C}^2/lattice$

and so in particular \tilde{B} is a Kähler variety. And that will done, by considering the local diffeomorphism

$$\mathbb{C}^2 \longrightarrow \tilde{B}, (\tau_1, \tau_2) \longmapsto \phi^{\tau_1} \phi^{\tau_2}(p),$$

for a fixed origin $p \in B$. The additive subgroup

$$\{(\tau_1, \tau_2) \in \mathbb{C}^2 : \phi^{\tau_1} \phi^{\tau_2}(p) = p\},$$

is a lattice of \mathbb{C}^2 , hence

$$\mathbb{C}^2 / \text{lattice} \longrightarrow \tilde{B},$$

is a biholomorphic diffeomorphism and \tilde{B} is a Kähler variety with Kähler metric given by $d\tau_1 \otimes d\bar{\tau}_1 + d\tau_2 \otimes d\bar{\tau}_2$. Now, a compact complex Kähler variety having the required number as (its dimension) of independent meromorphic functions is a projective variety [11]. In fact, here we have $\tilde{B} \subseteq \mathbb{P}^{15}$. Thus \tilde{B} is both a projective variety and a complex torus $\mathbb{C}^2 / \text{lattice}$ and hence an abelian surface as a consequence of Chow theorem. By the classification theory of ample line bundles on abelian varieties, $\tilde{B} \simeq \mathbb{C}^2 / L_\Omega$ with period lattice given by the columns of the matrix

$$\begin{pmatrix} \delta_1 & 0 & a & c \\ 0 & \delta_2 & c & b \end{pmatrix}, \quad \text{Im} \begin{pmatrix} a & c \\ c & b \end{pmatrix} > 0,$$

and

$$\delta_1 \delta_2 = g(\mathcal{H}_\varepsilon) - 1 = 1,$$

implying $\delta_1 = \delta_2 = 1$. Thus \tilde{B} is principally polarized and it is the jacobian of the hyperelliptic curve \mathcal{H}_ε . This completes the proof of the proposition.

Remark 4.1 *We have seen that the reflection σ on the affine variety B amounts to the flip*

$$\sigma : (z_1, z_2, z_3, z_4, z_5) \longmapsto (z_1, z_2, -z_3, -z_4, z_5),$$

changing the direction of the commuting vector fields. It can be extended to the $(-Id)$ -involution about the origin of \mathbb{C}^2 to the time flip $(t_1, t_2) \mapsto (-t_1, -t_2)$ on \tilde{B} , where t_1 and t_2 are the time coordinates of each of the flows X_{F_1} and X_{F_2} . The involution σ acts on the parameters of the Laurent solution (7) as follows

$$\sigma : (t, \alpha, \beta, \gamma, \theta, \varepsilon) \longmapsto (-t, -\alpha, -\beta, -\gamma, -\theta, -\varepsilon),$$

interchanges the curves $\mathcal{H}_{\varepsilon=\pm i}$ (8) and the linear space L can be split into a direct sum of even and odd functions. Geometrically, this involution interchanges \mathcal{H}_i and \mathcal{H}_{-i} , i.e., $\mathcal{H}_{-i} = \sigma \mathcal{H}_i$.

Remark 4.2 Consider on \tilde{B} the holomorphic 1-forms dt_1 and dt_2 defined by $dt_i(X_{F_j}) = \delta_{ij}$, where X_{F_1} and X_{F_2} are the vector fields generated respectively by F_1 and F_2 . Taking the differentials of $\zeta = 1/z_1$ and $\xi = z_1/z_2$ viewed as functions of t_1 and t_2 , using the vector fields and the Laurent series (7) and solving linearly for dt_1 and dt_2 , we obtain as expected the hyperelliptic holomorphic differentials

$$\begin{aligned}\omega_1 &= dt_1|_{\mathcal{H}_\varepsilon}, \\ &= \frac{1}{\Delta} \left(\frac{\partial \xi}{\partial t_2} d\zeta - \frac{\partial \zeta}{\partial t_2} d\xi \right) |_{\mathcal{H}_\varepsilon}, \\ &= \frac{\alpha d\alpha}{\sqrt{P(\alpha)}}, \\ \omega_2 &= dt_2|_{\mathcal{H}_\varepsilon}, \\ &= \frac{1}{\Delta} \left(-\frac{\partial \xi}{\partial t_1} d\zeta - \frac{\partial \zeta}{\partial t_1} d\xi \right) |_{\mathcal{H}_\varepsilon}, \\ &= \frac{\sqrt{2} d\alpha}{2\sqrt{P(\alpha)}},\end{aligned}$$

with

$$P(\alpha) \equiv 4\alpha^6 - 4a\alpha^4 - 4c_1\alpha^2 + 2\varepsilon\sqrt{2}c_2\alpha - c_3,$$

and

$$\Delta \equiv \frac{\partial \zeta}{\partial t_1} \frac{\partial \xi}{\partial t_2} - \frac{\partial \zeta}{\partial t_2} \frac{\partial \xi}{\partial t_1}.$$

The zeroes of ω_2 provide the points of tangency of the vector field X_{F_1} to \mathcal{H}_ε . We have $\frac{\omega_1}{\omega_2} = -\varepsilon\sqrt{2}\alpha$, and X_{F_1} is (doubly) tangent to \mathcal{H}_ε at the point covering $\alpha = \infty$, i.e., where both the curves touch.

References

- [1] *M. Adler, P. van Moerbeke.* The complex geometry of the Kowalewski-Painlevé analysis. *Invent. Math.* 97 (1989) 3-51.
- [2] *A.I. Belokolos, V.Z. Bobenko, V.Z. Enol'skii, A.R. Its, V.B. Matveev.* *Algebro-Geometric approach to nonlinear integrable equations.* Springer-Verlag 1994.
- [3] *P.A. Griffiths, J. Harris.* *Principles of algebraic geometry.* Wiley-Interscience 1978.
- [4] *L. Haine.* Geodesic flow on $SO(4)$ and Abelian surfaces. *Math. Ann.* 263 (1983) 435-472.
- [5] *A. Lesfari.* Abelian surfaces and Kowalewski's top. *Ann. Scient. École Norm. Sup. Paris sér. 4*, 21 (1988) 193-223.

- [6] *A. Lesfari*. Completely integrable systems : Jacobi's heritage. J. Geom. Phys. 31 (1999) 265-286.
- [7] *A. Lesfari*. Le théorème d'Arnold-Liouville et ses conséquences. Elem. Math. 58 (2003) 6-20.
- [8] *A. Lesfari*. Analyse des singularités de quelques systèmes intégrables. C.R. Acad. Sci. Paris, Ser. I, 341 (2005) 85-88.
- [9] *A. Lesfari*. Abelian varieties, surfaces of general type and integrable systems. Beiträge Algebra Geom., Vol.48, 1 (2007) 95-114.
- [10] *B.G. Moishezon*. On n -dimensional compact varieties with n algebraically independent meromorphic functions. Amer. Math. Soc. Transl. 63 (1967) 51-177.
- [11] *D. Mumford*. On the equations defining abelian varieties I, II, III. Invent. Math. 1 (1966) 287-354 ; 3 (1967) 75-135; 3 (1967) 215-244.