

The Hénon-Heiles system as part of an integrable system in five unknowns with three constants of motions

A. Lesfari

*Department of Mathematics
Faculty of Sciences
University of Chouaïb Doukkali
B.P. 20, El Jadida, Morocco.
Lesfariahmed@yahoo.fr*

Abstract. In this paper we construct a new completely integrable system. This system is an instance of a master system of differential equations in 5 unknowns having 3 quartics constants of motion. We find via the Painlevé analysis the principal balances of the hamiltonian field defined by the hamiltonian. Consequently, the system in question is algebraically integrable. A careful analysis of this system reveals an intimate rational relationship with a special case of the well known Hénon-Heiles system. The latter admits asymptotic solutions with fractional powers in t and depending on 3 free parameters. As a consequence, this system is algebraically completely integrable in the generalized sense.

2010 Mathematics Subject Classification : 70H06, 37J35.

Key words and phrases : Hamiltonian, Integrable systems.

Among recent researches, there have been many works dealing with the integrability of hamiltonian systems. In hamiltonian mechanics, for integrating a dynamical system with n degrees of freedom, it is sufficient in most cases to know only the first n integrals. This situation is known as Liouville complete integrability of a hamiltonian system. A dynamical system is algebraic completely integrable (in the sense of Adler-van Moerbeke [1,2]) if it can be linearized on a complex algebraic torus $\mathbb{C}^n/lattice$ (=abelian variety). The invariants (often called first integrals or constants) of the motion are polynomials and the phase space coordinates (or some algebraic functions of these) restricted to a complex invariant variety defined by putting these invariants equals to generic constants, are meromorphic functions on an abelian variety. Moreover, in the coordinates of this abelian variety, the flows (run with complex time) generated by the constants of the motion are straight lines. Also some interesting integrable systems appear as cov-

erings of algebraic completely integrable systems [4,9,12,14]. The invariant varieties are coverings of abelian varieties and these systems are called algebraic completely integrable in the generalized sense. Beside the fact that many integrable hamiltonian systems have been on the subject of powerful and beautiful theories of mathematics, another motivation for its study is : the concepts of integrability have been applied to an increasing number of physical systems, biological phenomena, population dynamics, chemical rate equations, to mention only a few. However, we have no general method for testing the integrability of a given hamiltonian system. It seems still hopeless to describe, or even to recognize with any facility, those hamiltonian systems which are integrable, though they are quite exceptional. In the present paper, we present an integrable system in five unknowns having two cubic and one quartic invariants (constants of motion). Also, we show that this system include in particular the well known Hénon-Heiles system.

Consider the system

$$\begin{aligned}\dot{y}_1 &= x_1, \\ \dot{y}_2 &= x_2, \\ \dot{x}_1 &= -Ay_1 - 2y_1y_2, \\ \dot{x}_2 &= -By_2 - y_1^2 - \varepsilon y_2^2,\end{aligned}\tag{1}$$

corresponding to a generalized Hénon-Heiles hamiltonian

$$H = \frac{1}{2}(x_1^2 + x_2^2) + \frac{1}{2}(Ay_1^2 + By_2^2) + y_1^2y_2 + \frac{\varepsilon}{3}y_2^3,$$

where A, B, ε are constant parameters and y_1, y_2, x_1, x_2 are canonical coordinates and momenta, respectively. First studied as a mathematical model to describe the chaotic motion of a test star in an axisymmetric galactic mean gravitational field [7], this system is widely explored in other branches of physics. It well-known from applications in stellar dynamics, statistical mechanics and quantum mechanics. It provides a model for the oscillations of atoms in a three-atomic molecule [5].

Usually, the Hénon-Heiles system is not integrable and represents a classical example of chaotic behaviour. Nevertheless at some special values of the parameters it is integrable; to be precise, there are known three integrables cases :

(i) $\varepsilon = 6$, A and B arbitrary. The second integral of motion is

$$H_2 = y_1^4 + 4y_1^2y_2^2 - 4x_1^2y_2 + 4x_1x_2y_1 + 4Ay_1^2y_2 + (4A - B)x_1^2 + A(4A - B)y_1^2.$$

(ii) $\varepsilon = 1$, $A = B$. The second integral of motion is

$$H_2 = x_1x_2 + \frac{1}{3}y_1^3 + y_1y_2^2 + Ay_1y_2.$$

(iii) $\varepsilon = 16$, $B = 16A$. The second integral of motion is

$$H_2 = 3x_1^4 + 6Ax_1^2y_1^2 + 12x_1^2y_1^2y_2 - 4x_1x_2y_1^3 - 4Ay_1^4y_2 - 4y_1^4y_2^2 + 3A^2y_1^4 - \frac{2}{3}y_1^6. \quad (2)$$

In the two cases (i) and (ii), the system (1) has been integrated by making use of genus one and genus two theta functions. For the case (i), it was shown [3] that this case separates in translated parabolic coordinates. Solving the problem in case (ii) is not difficult (this case trivially separates in cartesian coordinates). In the case (iii), the system can also be integrated [13] by making use of elliptic functions.

The general solutions of the equations of motion for hamiltonian (1), for the case (i) and (ii), have the Painlevé propriety, i.e., that they admit only poles in the complex time variable. For the case (ii), we found [8,10,11] via Kowalewski-Painlevé analysis the principal balances of the hamiltonian vector field defined by the hamiltonian and we have shown that the system is algebraic complete integrable. The affine surface defined by the two constants of motion completes into an abelian surface by adjoining a smooth genus three hyperelliptic curve D . This curve is a double cover of an elliptic curve E , ramified at four points : it defines a line bundle and a polarization (1,2) on this abelian surface. Also, the abelian surface obtained is the dual of a Prym variety, namely $\text{Prym}(D/E)^\vee$ and the flow of solutions of the equations of motion is linearized on abelian surface.

The present paper deals with the case (iii). The system (1) can be written in the form

$$\dot{u} = J \frac{\partial H}{\partial u}, \quad u = (y_1, y_2, x_1, x_2)^\top, \quad (3)$$

where

$$H \equiv H_1 = \frac{1}{2}(x_1^2 + x_2^2) + \frac{A}{2}(y_1^2 + 16y_2^2) + y_1^2y_2 + \frac{16}{3}y_2^3, \quad (4)$$

$$\frac{\partial H}{\partial z} = \left(\frac{\partial H}{\partial y_1}, \frac{\partial H}{\partial y_2}, \frac{\partial H}{\partial x_1}, \frac{\partial H}{\partial x_2} \right)^\top, \quad J = \begin{pmatrix} O & I \\ -I & O \end{pmatrix}.$$

The functions $H_1(4)$ and $H_2(2)$ commute :

$$\{H_1, H_2\} = \sum_{k=1}^2 \left(\frac{\partial H_1}{\partial p_k} \frac{\partial H_2}{\partial q_k} - \frac{\partial H_1}{\partial q_k} \frac{\partial H_2}{\partial p_k} \right) = 0.$$

The second flow commuting with the first is regulated by the equations

$$\dot{u} = J \frac{\partial H_2}{\partial u}, \quad u = (y_1, y_2, x_1, x_2)^\top$$

and is written explicitly as

$$\begin{aligned} \dot{y}_1 &= -24Ax_1 - 8x_1y_2 + 4x_2y_1, \\ \dot{y}_2 &= 4x_1y_1, \\ \dot{x}_1 &= 24A^2y_1 - 4x_1x_2 - 8Ay_1y_2 - 8y_1y_2^2 - 4y_1^3, \\ \dot{x}_2 &= 4x_1^2 - 4Ay_1^2 - 8y_1^2y_2, \end{aligned}$$

The invariant (or level) variety

$$A = \bigcap_{k=1}^2 \{z \in \mathbb{C}^4 : H_k(z) = b_k\}, \quad (5)$$

is a smooth affine surface for generic $(b_1, b_2) \in \mathbb{C}^2$. It was conjectured in [13] that the complex invariant variety $A(5)$ is an affine part of an abelian variety (an algebraic complex torus). We will see that in fact the invariant surface $A(5)$ can be completed as a cyclic double cover \overline{A} of an abelian surface \widetilde{B} . The system (3) is algebraic complete integrable in the generalized sense. Moreover, \overline{A} is smooth except at the point lying over the singularity of type $A_1 : X^2 + Y^2 - Z^2 = 0$ and the resolution \hat{A} of \overline{A} is a surface of general type.

When one examines all possible singularities, one finds that it possible for the variable y_1 to contain square root terms of the type $t^{1/2}$, which are strictly not allowed by the Painlevé test. However, these terms are trivially removed by introducing some new variables z_1, \dots, z_5 (see below), which restores the Painlevé property to the system. The system (3) admits Laurent solutions in $t^{1/2}$, depending on three free parameters: α, β, γ and they are explicitly given as follows

$$\begin{aligned} y_1 &= \frac{\alpha}{\sqrt{t}} + \beta t \sqrt{t} - \frac{\alpha}{18} t^2 \sqrt{t} + \frac{\alpha A^2}{10} t^3 \sqrt{t} - \frac{\alpha^2 \beta}{18} t^4 \sqrt{t} + \dots, \\ y_2 &= -\frac{3}{8t^2} - \frac{A}{2} + \frac{\alpha^2}{12} t - \frac{2A^2}{5} t^2 + \frac{\alpha\beta}{3} t^3 - \gamma t^4 + \dots, \\ x_1 &= -\frac{1}{2} \frac{\alpha}{t\sqrt{t}} + \frac{3}{2} \beta \sqrt{t} - \frac{5}{36} \alpha t \sqrt{t} + \frac{7}{20} \alpha A^2 t^2 \sqrt{t} - \frac{1}{4} \alpha^2 \beta t^3 \sqrt{t} + \dots, \\ x_2 &= \frac{3}{4t^3} + \frac{1}{12} \alpha^2 - \frac{4}{5} A^2 t + \alpha \beta t^2 - 4\gamma t^3 + \dots \end{aligned} \quad (6)$$

These formal series solutions are convergent as a consequence of the majorant method. By substituting these series in the constants of the motion $H_1 = b_1$ and $H_2 = b_2$, i.e.,

$$\begin{aligned} H_1 &= \frac{1}{9} \alpha^2 - \frac{21}{4} \gamma + \frac{13}{288} \alpha^4 + \frac{4}{3} A^3 = b_1, \\ H_2 &= -144 \alpha \beta^3 + \frac{294}{5} \alpha^3 \beta A^2 + \frac{8}{9} \alpha^6 - 33 \gamma \alpha^4 = b_2, \end{aligned}$$

one eliminates the parameter γ linearly, leading to an equation connecting the two remaining parameters α and β :

$$144 \alpha \beta^3 - \frac{294 A^2}{5} \alpha^3 \beta + \frac{143}{504} \alpha^8 - \frac{4}{21} \alpha^6 + \frac{44}{21} (4A^3 - 3b_1) \alpha^4 + b_2 = 0.$$

which is nothing but the equation of an algebraic curve \mathcal{D} of genus four along which the $u(t) \equiv (y_1(t), y_2(t), x_1(t), x_2(t))$ blow up. To be more precise \mathcal{D} is

the closure of the continuous components of

$$\{\text{Laurent series solutions } u(t) \text{ such that } H_k(u(t)) = b_k, 1 \leq k \leq 2\},$$

i.e.,

$$\mathcal{D} = t^0\text{-coefficient of } \{u \in \mathbb{C}^4 : H_1(u(t)) = b_1\} \cap \{u \in \mathbb{C}^4 : H_2(u(t)) = b_2\}.$$

The Laurent solutions restricted to the surface $A(5)$ are parameterized by this curve \mathcal{D} .

We show that the system (3) is part of a new system of differential equations in five unknowns having two cubic and one quartic invariants (constants of motion). By inspection of the expansions (6), we look for polynomials in (y_1, y_2, x_1, x_2) with a simple pole. Let

$$\varphi : A \longrightarrow \mathbb{C}^5, (y_1, y_2, x_1, x_2) \longmapsto (z_1, z_2, z_3, z_4, z_5), \quad (7)$$

be a morphism on the affine variety $A(5)$ where z_1, \dots, z_5 are defined as

$$z_1 = y_1^2, \quad z_2 = y_2, \quad z_3 = x_2, \quad z_4 = y_1 x_1, \quad z_5 = 3x_1^2 + 2y_1^2 y_2.$$

The morphism (7) maps the vector field (3) into a Liouville integrable system (8) in five unknowns having two cubic and one quartic invariants. Indeed, the change of variables (7) maps the vector field (4) into the system of differential equation

$$\begin{aligned} \dot{z}_1 &= 2z_4, \\ \dot{z}_2 &= z_3, \\ \dot{z}_3 &= -z_1 - 16Az_2 - 16z_2^2, \\ \dot{z}_4 &= -Az_1 + \frac{1}{3}z_5 - \frac{8}{3}z_1 z_2, \\ \dot{z}_5 &= -6Az_4 + 2z_1 z_3 - 8z_2 z_4, \end{aligned} \quad (8)$$

with constants of motion

$$\begin{aligned} F_1 &= \frac{1}{2}Az_1 + \frac{1}{6}z_5 + 8Az_2^2 + \frac{1}{2}z_3^2 + \frac{2}{3}z_1 z_2 + \frac{16}{3}z_2^3, \\ F_2 &= 9A^2 z_1^2 + z_5^2 + 6Az_1 z_5 - 2z_1^3 - 24Az_1^2 z_2 - 12z_1 z_3 z_4 + 24z_2 z_4^2 - 16z_1^2 z_2^2, \\ F_3 &= z_1 z_5 - 3z_4^2 - 2z_1^2 z_2. \end{aligned}$$

This new system is completely integrable and the hamiltonian structure is defined by the Poisson bracket

$$\{F, H\} = \left\langle \frac{\partial F}{\partial z}, J \frac{\partial H}{\partial z} \right\rangle = \sum_{k,l=1}^5 J_{kl} \frac{\partial F}{\partial z_k} \frac{\partial H}{\partial z_l},$$

where

$$\frac{\partial H}{\partial z} = \left(\frac{\partial H}{\partial z_1}, \frac{\partial H}{\partial z_2}, \frac{\partial H}{\partial z_3}, \frac{\partial H}{\partial z_4}, \frac{\partial H}{\partial z_5} \right)^\top,$$

and

$$J = \begin{pmatrix} 0 & 0 & 0 & 2z_1 & 12z_4 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & -1 & 0 & 0 & -2z_1 \\ -2z_1 & 0 & 0 & 0 & -8z_1z_2 + 2z_5 \\ -12z_4 & 0 & 2z_1 & 8z_1z_2 - 2z_5 & 0 \end{pmatrix},$$

is a skew-symmetric matrix for which the corresponding Poisson bracket satisfies the Jacobi identities. The system (8) can be written as

$$\dot{z} = J \frac{\partial H}{\partial z}, \quad z = (z_1, z_2, z_3, z_4, z_5)^\top,$$

where $H = F_1$. The second flow commuting with the first is regulated by the equations

$$\dot{z} = J \frac{\partial F_2}{\partial z}, \quad z = (z_1, z_2, z_3, z_4, z_5)^\top,$$

and is written explicitly as

$$\begin{aligned} \dot{z}_1 &= 24z_4z_5 - 24z_1^2z_3 + 96z_1z_2z_4 + 72Az_1z_4, \\ \dot{z}_2 &= -12z_1z_4, \\ \dot{z}_3 &= 12Az_1^2 - 12z_1z_5 + 48z_1^2z_2, \\ \dot{z}_4 &= 4z_5^2 - 36A^2z_1^2 + 12z_1^3 - 16z_1z_2z_5 + 48Az_1^2z_2 + 24z_1z_3z_4 + 64z_1^2z_2^2, \\ \dot{z}_5 &= -96z_2z_4z_5 + 768z_1z_2^2z_4 - 72Az_4z_5 + 576Az_1z_2z_4 + 144z_3z_4^2 \\ &\quad - 216A^2z_1z_4 + 48z_1^2z_4 - 96z_1^2z_2z_3 + 24z_1z_3z_5. \end{aligned}$$

These vector fields are in involution, i.e.,

$$\{F_1, F_2\} = \left\langle \frac{\partial F_1}{\partial z}, J \frac{\partial F_2}{\partial z} \right\rangle = 0,$$

and the remaining one is casimir, i.e.,

$$J \frac{\partial F_3}{\partial z} = 0.$$

Consequently, the system (8) is integrable in the sense of Liouville.

The invariant variety

$$B = \bigcap_{k=1}^3 \{z \in \mathbb{C}^5 : F_k(z) = c_k\}, \quad (9)$$

is a smooth affine surface for generic values of c_1, c_2 and c_3 . The system (8) possesses Laurent series solutions which depend on four free parameters.

These meromorphic solutions restricted to the surface $B(9)$ can be read off from (6) and the change of variable (7). Following the methods in [1,2,11], one find the compactification of B into an abelian surface \tilde{B} , the system of differential equations (8) is algebraic complete integrable and the corresponding flows evolve on \tilde{B} . Applying the method explained in Piovan [20] (see also [9]), we show that the invariant surface $A(5)$ can be completed as a cyclic double cover \overline{A} of an abelian surface \tilde{B} . The system (3) is algebraic complete integrable in the generalized sense. Moreover, \overline{A} is smooth except at the point lying over the singularity of type $A_1 : X^2 + Y^2 - Z^2 = 0$ and the resolution \tilde{A} of \overline{A} is a surface of general type. We have shown that the morphism φ (9) maps the vector field (4) into an algebraic completely integrable system (8) in five unknowns and the affine variety A (5) onto the affine part B (9) of an abelian variety \tilde{B} . This explains (among other) why the asymptotic solutions to the differential equations (3) contain fractional powers.

In this paper we have shown that the Hénon-Heiles (third case) equations are part of a system of differential equations in five unknowns having three constants of motions. We have seen that this new system is Liouville integrable but we do not know physical interpretation for this system and it may be possible to have links with other systems (in addition to Hénon-Heiles) with applications in physics among other. The integrable dynamical system presented here is an interesting example, particular to experts of abelian varieties who may want to see an explicit example of a correspondence for varieties defined by different curves.

References

- [1] Adler, M. and van Moerbeke, P.: The complex geometry of the Kowalewski-Painlevé analysis. *Invent. Math.*, 97, 3-51 (1989).
- [2] Adler, M., van Moerbeke, P. and Vanhaecke, P.: Algebraic integrability, Painlevé geometry and Lie algebras. A series of modern surveys in mathematics, Volume 47, Springer-Verlag, 2004.
- [3] Ankiewicz, A. and Pask, C.: The Complete Whittaker Theorem For Two-Dimensional Integrable Systems And Its Application. *J Phys A-Math Gen*, 16: 4203-4208 (1983).
- [4] Bechlivanidis, C., van Moerbeke, P.: The Goryachev Chaplygin top, the Toda lattice and the intersection of 5 quadrics in \mathbb{P}^7 . *Commun. Math. Phys.*, 10, 317-324 (1987).
- [5] Berry, M.L.: Regular and irregular motions. Topics in nonlinear dynamics, S. Jorna (ed.), *Am. Inst. Phys.*, 16-120, New York 1978.

- [6] Griffiths, P.A., Harris,J.: Principles of algebraic geometry. Wiley-Interscience 1978.
- [7] Hénon, M. and Heiles, C.: The applicability of the third integral of motion; some numerical experiments. *Astron. J.*, 69, 73-79 (1964).
- [8] Lesfari, A. : Le système différentiel de Hénon-Heiles et les variétés Prym. *Pacific J. Math.*, Vol. 212, N°1, 125-132 (2003).
- [9] Lesfari, A.: Abelian varieties, surfaces of general type and integrable systems. *Beiträge Algebra Geom.*, Vol.48, 1, 95-114 (2007).
- [10] Lesfari, A.: Integrable systems and complex geometry. *Lobachevskii Journal of Mathematics*, Vol.30, No.4, 292-326 (2009).
- [11] Lesfari, A.: Algebraic integrability : the Adler-van Moerbeke approach. *Regul. Chaotic Dyn.*, Vol. 16, Nos.3-4, pp.187-209 (2011).
- [12] Piovan, L.: Cyclic coverings of abelian varieties and the Goryachev-Chaplygin top. *Math. Ann.*, 294, 755-764 (1992).
- [13] V. Ravoson, L. Gavrilov, and R. Caboz, Separability and Lax pairs for Hénon-Heiles system. *J. Math. Phys.*, 34, 2385-2393 (1993).
- [14] Vanhaecke, P., Integrable Systems in the Realm of Algebraic Geometry, 2nd ed., Lecture Notes in Math., vol. 1638, Berlin: Springer, 2001.